

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## CURRENT LITERATURE

## MINOR NOTICES

Submerged forests.—People invariably are fascinated by evidences of former forests below present sea-levels, as indicated by the appearance at low tide of stumps in situ. That these "Noah's woods" are of large scientific interest is well brought out in an interesting little booklet by Clement Reid, forming one of the useful series of "Cambridge Manuals of Science and Literature." Reid speaks in detail of the various submerged forests of the English coast and of the Dogger Banks in the North Sea. The lowest of these forests are oak forests much like those of the present day, though apparently with fewer species. Among the species absent are the more southern types that have come into England since the ice age; at the same time, the glacial relicts have disappeared. It is concluded that the lowest of these forests date back about 5000 years, and that there has been no essential change of sea-level in the last 3500 years. During the period of the life of these forests the climate was much like that of the present, and the land was 60–90 feet higher than it is today.

In connection with Reid's book on submerged forests, attention may be called to a short paper by Osborn, dealing with a submerged forest on the coast of Wales.<sup>2</sup>—H. C. Cowles.

Plant evolution.—Campbell's has presented an interesting popular account of plant evolution for the "American Nature Series." Following a brief consideration of the factors in evolution, the different plant groups are considered, from the bacteria and blue-green algae through the angiosperms, the object here being to set forth the chief characters of each group in the scale of ascent. The book closes with four chapters having the following titles: environment and adaptation, the problems of plant distribution, the human factor in evolution, and the origin of species. Since most books of this character deal with animals rather than with plants, this volume of Campbell's is especially welcome.—H. C. Cowles.

<sup>&</sup>lt;sup>1</sup> Reid, Clement, Submerged forests. pp. 129. figs. 4 and frontispiece. Cambridge University Press. 1913.

<sup>&</sup>lt;sup>2</sup> OSBORN, T. G. B., A note on the submerged forest at Llanaber, Barmouth. pp. 10. figs. 2. pls. 2. Reprint from Mem. Proc. Manchester Lit. Phil. Soc. 56: 1912.

<sup>&</sup>lt;sup>3</sup> CAMPBELL, D. H., Plant life and evolution. 8vo. pp. iv+36o. figs. 22. New York: Henry Holt & Co. 1911. \$1.60.